

GEL'FENBEIN, S. Kh.

Author: Gel'fenbein, S. Kh.

Title: The Gas Turbine in domestic industry. (Gazovye turbiny; bibliografiia. 23 p.

City: Moscow

Publisher:

~~Publishing House~~ State Printing House of the Machine Construction literature.

Date: 1946

Available: Library of Congress

Source: Monthly List of Russian Accessions, V. 3, No. 12, March 1951

GEL'FENBERG, S. KH.

Gazovaia turbina v domennom proizvodstve. Moskva, Mashgiz, 1946. 29, (3) p. illus.

The gas turbine in blast-furnace production.

DLC: TN673.G4

SO: Manufacturing and Mechanical Engineering in the Soviet Union, Library of Congress, 1953.

GEL'FENBEYN, S. KH

Aug 1947

USSR/Engineering  
Turbines, Gas  
Turbines - Combustion

"Experimental Research and Complex Gas Turbine Plant  
with Constant Pressure Combustion," C. Kh. Gel'fen-  
beyn, Candidate in Technical Sciences, Central Sci-  
entific Research Boiler Turbine Institute issled-  
I. I. Polzunov, 1 p

"Kotloturbostroeniye" No 4

A brief preliminary report on the outcome of experi-  
mental research done at the Laboratory of Gas Tur-  
bines TSERL on gas turbine plants (p-constant),  
carried out in the simplest plan with high-speed

24726

Aug 1947

USSR/Engineering (Contd)

axial compressor, gas turbines, and chamber combus-  
tion. As a result of the tests, they acquired a  
series of valuable methods for measuring discharge,  
and for measuring pressure and temperature during  
comparatively great speeds in axial clearances of  
near axial compressors and gas turbines.

24726

GEL'FERBEIN, S. KE.

Heat of dilution of sulfuric-nitric acid mixtures on the  
heat of reaction of these acids. M. S. G. (Sov. Chem. Soc. 1950, 20, 113-19)  
(Sov. Chem. Soc. 1950, 20, 113-19)  
(Sov. Chem. Soc. 1950, 20, 113-19)  
(1950).—The heat of reaction  $Q$  of  $\text{HNO}_3$  with  $\text{H}_2\text{SO}_4$  de-  
creases with increasing diln. If  $Q$  of the pure acids is taken  
= 1, the heat evolved in the interaction of the acids taken  
in the same proportions, but diln. with  $\text{H}_2\text{O}$ , is a fraction  
of  $Q$  of the pure acids; the decrease of  $Q$  with increasing  $\text{H}_2\text{O}$   
content of the mixt. is nearly linear. In terms of the pro-  
portions of  $\text{HNO}_3$  and  $\text{H}_2\text{SO}_4$  in the undiln. mixt.,  $Q$  passes  
through a max. (at about 67 wt. %  $\text{H}_2\text{SO}_4$ ). The  $Q$  plots  
permit calcn. of  $Q$  for any mixt. at any diln. The accepted  
representation of the reaction is  $\text{HNO}_3 + \text{H}_2\text{SO}_4 = \text{NO}_3^- +$   
 $\text{HSO}_4^- + \text{H}_2\text{O}$ . This scheme accounts for the high elec.  
cood. of the mixt. of the undiln. acids, and (in contrast to  
the behavior of the single acids) its fall with increasing diln.  
which shifts the equil. to the left and thus decreases the  
concen. of ions. This effect accounts also for the decrease  
of  $Q$  with increasing diln. In concert calcn. of heats of  
nitration, two factors, not usually taken into consideration,  
should be allowed for. One is the heat of sepn. of  $\text{HNO}_3$   
and  $\text{H}_2\text{SO}_4$  in the nitration mixt.; this is equal to the heat of  
mixing, with reversed sign. The other factor is the heat of  
diln. not of the single  $\text{H}_2\text{SO}_4$ , but of the  $\text{H}_2\text{SO}_4 + \text{HNO}_3$   
mixt. (with 20–40% excess  $\text{HNO}_3$ ) commonly used in nitra-  
tion; this heat of diln. of the mixt. of acids can be evaluated  
from the difference of the heat of formation of the aq. mixt.  
and the heat of mixing of the undiln. acids. As ex-  
amples, detailed calcns. are given of heats of nitration of  
glycerol and of  $\text{PhOH}$ . With 1 ton of glycerol, and 5000  
kg. of acid ( $\text{HNO}_3$  50,  $\text{H}_2\text{SO}_4$  50%), the resulting acid is  
composed of  $\text{HNO}_3$  7.45,  $\text{H}_2\text{SO}_4$  25.5,  $\text{H}_2\text{O}$  22.53 kilograms;  
the total heat evolved is 251,669 kcal. as against 347,000  
kcal. calcd. by the usual procedure, i.e. 27% too high.  
Similarly, for the nitration of  $\text{PhOH}$ , the conventional  
method gives a heat by 31% too high. The discrepancy  
becomes even more serious (89% for glycerol, 25% for  
 $\text{PhOH}$ ) if experimentally detd. heats of diln. are taken in-  
stead of the data of McDavid (C. 4, 16, 3733). The new  
exptl. detns. give (g. mixt., initial compn. in %  $\text{HNO}_3$ ,  
 $\text{H}_2\text{SO}_4$ ,  $\text{H}_2\text{O}$ , g.  $\text{H}_2\text{O}$  added, final compn. in %  $\text{HNO}_3$ ,  
 $\text{H}_2\text{SO}_4$ ,  $\text{H}_2\text{O}$ , heat of diln. in cal.): 45.0, 15.2, 85.1, —, 2.3,  
14.4, 80.75, 4.85, 485; 47.3, 14.4, 80.75, 4.85, 3.7, 13.4,  
75.35, 11.25, 495; 51.05, 13.4, 75.35, 11.25, 2.84, 12.70,  
70.85, 16.45, 298; 50.0, 2.53, 88.68, 8.8, 3.5, 2.3, 82.9,  
14.8, 516; 45.0, 9.18, 84.53, 8.24, 2.7, 8.4, 80.1, 11.5, 305.  
N. Thon

GEL'FENBYN, S.Kh., kand.tekhn.nauk

Alternating working conditions of a gas turbine. Sudostroenie  
no.7:23-29 J1 '60. (MIRA 13:7)  
(Marine gas turbines)

S/262/62/000/008/010/022  
I007/I207

AUTHOR: Gel'fenbeyn, S. Kh.

TITLE: Method of computing variable operating conditions for marine gas turbines

PERIODICAL: Referativnyy zhurnal, otdel'nyy vypusk. 42. Silovyye ustanovki, no. 8, 1962, 35, abstract 42.8.182. "Tr. Tsent. n.-i. in-ta morsk. flota", no. 34, 1961, 24-38

TEXT: It is shown that operation of the compressor is approximately proportional to the square of its rotational speed. A formula is derived for computing the relationship between gas consumption and rotational speed of the turbine driving the compressor in the single-shaft and double-shaft gas turbine unit. The formula was obtained analytically utilizing empirical relationships.



[Abstracter's note: Complete translation.]

Card 1/1

GEL'FENBEYN, S.Kh., kand.tekhn.nauk

Rules of the U.S.S.R. Register on marine gas turbine plants.  
Sudostroenie 28 no.3:21-23 Mr '62. (MIRA 15:4)  
(Marine gas turbines) (Ship registers)

GEL'FENBEYN, S.Kh., kand.tekhn.nauk, dotsent

Calculation of the regenerator of a gas turbine system for operation on principal parameters and partial load conditions. Energomashinostroenie 9 no.11:25-29 N '63. (MIRA 17:2)



L 11230-63

EPA/EWT(m)/BDS--AEDC/AFFTC/ASD--Paa-4

ACCESSION NR: AP3001476

S/0114/63/000/006/0037/0038

57

AUTHOR: Gel'fenbeyn, S. Kh. (Candidate of technical sciences)

TITLE: Testing the gas turbine with a free-piston gas producer

SOURCE: <sup>21</sup> Energomashinostroyeniye, no. 6, 1963, 37-38

TOPIC TAGS: gas turbine, free-piston gas producer, "Pavlin Vinogradov" ship

ABSTRACT: Results are reported of testing in 1961 a six-stage gas turbine and its free-piston gas producer as a main propulsion plant on the "Pavlin Vinogradov" ship. The tests were carried out under mooring, ballast, and cargo conditions. The turbine plant was built by the "Sigma" Company. The techniques and results of measurements are reported in detail. The turbine efficiency (under ballast conditions) in the power range 34-100 per cent, or rpm range 74-100 per cent, varied within 9 per cent. In stand tests the efficiency varied little within 23.5-100 per cent power, or 62-100 per cent rpm. Orig. art. has: 3 figures and 2 formulas.

ASSOCIATION: none

SUBMITTED: 00

DATE ACQD: 01Jul63

ENCL: 00

SUB CODE: 00

NO REF SOV: 002

OTHER: 000

Cord 1/1 ch/wm

GEI FENBEYN, S.M., kand.takhl.nark, dozent

Heat exchangers for closed-cycle type gas turbine engines.  
Energomashinostroitel' no. 3: 22-33 Ag '65.

(MIRA 16:10)

DUBROVSKIY, A.A., kand. tekhn.nauk; GEL'FENBEYN, S.P., inzh.

Using negative feedback for improving the efficiency of machinery.  
Mekh. i elek. sots. sel'khoz. 19 no.3:30-32'61. (MIRA 4:6)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut mekhanizatsii  
sel'skogo khozyaystva.  
(Agricultural machinery)

DUBROVSKIY, A. A., kand. tekhn. nauk; GEL'FENBEYN, S. P., inzh.

Some problems concerning the automation of mobile agricultural  
machines. Mekh. i elek. sots. sel'khoz. 20 no.6:21-25 '62.  
(MIRA 16:1)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut mekhanizatsii  
sel'skogo khozyaystva.

(Automation) (Farm mechanization)

CEL'FENBEYN, S.P., kand. tekhn. nauk; SVIRSHCHEVSKIY, A.V., kand. tekhn. nauk

Conference on the Automation of Agriculture. Mekh. i elek. sots.  
sel'khoz. 21 no.5:36,58 '63. (MIRA 17:1)

GEL'FENBYN, YA. V.

15/49T72

USSR/Engineering  
Pumps, Plunger  
Machines, Testing

Aug 48

"Mechanical Pressure Testing of Boilers, Pipes and  
Armatures," Ya. V. Gel'fenbeyn, Engr,  $\frac{1}{2}$  p

"Mekh Stroi" No 8

Describes test pump produced by Glavpromenergomontazh  
Plant in Leningrad. It is horizontal plunger type,  
driven by an electric motor through worm reduction  
gearing. Maximum pressure 150 ats.

15/49T72

GEL'FENBEYN, Ya. V.

20657 Gel'fenbeyn. Ya. V. Mekhanizatsiya priteski i Rezki Kirpicha. Mekhanizatsiya  
stroit-va, 1949, No. 6, s. 22-23

SO: LETOPIS ZHURNAL STATEY - Vol. 28, Moskva, 1949

GEL'FENBEYN, Ya. V., Eng.

Devices used for installation work on boiler units and plant pipe lines.  
Rab. energ., 1, No 2, 1951.



~~Feas.~~ Gel'fenbeyn Ya. V.

AID P - 4062

Subject : USSR/Power  
Card 1/1 Pub. 26 - 20/33  
Authors : Alekseyev, N. V. and Ya. V. Gel'fenbeyn, Engs.  
Title : Applying plastering solutions without using a compressor.  
Periodical : Elek. sta., 12, 49-50, 1955  
Abstract : Various methods of applying plaster without using a compressor but depending upon the size and quality of sand used are explained. Three diagrams.  
Institution : None  
Submitted : No date

GEL'FENBEYN, Ya.V., inzh.

Inflated temporary structures for the construction of thermal electric plants. Energ. stroi. no.22:24-25 '61. (MIRA 15:7)

1. Proyechnaya tresta "Sevenenergoprojekt".  
(Buildings, Portable) (Electric power plants)

GEL'FENBEYN, Ya. V., inzh.

New truck-mounted cranes abroad. Stro1. 1 dor. mash. 8 no.5:  
33-35 My '63. (MIRA 16:5)  
(Cranes, derricks, etc.)

GEL'FENBEYN, Ya.V., inzh.

Installing tracks for gantry cranes. Prom. stroi. 40 [i.e. 41],  
no.5:47-48 My '63. (MIRA 16:5)  
(Cranes, derricks, etc.)

GEL'FENBEYN, Ya.V., irzh.

Crawler and automobile cranes in foreign countries. Stroil. 1  
dor. mash. 9 no.9:34-38 S '64. (MIRA 17:11)

CEL'FANDHEYN, Ya. V. (Page)

Determination of the dynamic characteristics of a closed-loop  
nonstationary system in normal operation. Izv. AN SSSR. Tekh.  
kib. no.4:111-114 J1-kg '65. (MIRA 18:11)

GEL'FENBEYN, Ya.V., inzh.

Need for the publishing of a "Manual on the preparation of organization plans for the construction of thermal electric power plants." Energ.stroi. no.25:94-96 '61. (MIRA 15:4)  
(Electric power plants--Design and construction)

GEL'FENBEYN, Yakov Vladimirovich; VINITSKIY, D.Ya., inzh., retsenzent;  
EBERLIN, L.A., inzh., retsenzent; PAVLOV, N.G., red.;  
ZHITNIKOVA, O.S., tekhn. red.

[Hoisting mechanisms and rigging devices for the construction  
of thermal electric power plants] Gruzopod'emnye mekhanizmy i  
takelazhnye prispособleniia dlia stroitel'stva teplovykh  
elektrostantsii. Moskva, Gosenergoizdat, 1962. 270 p.

(MIRA 15:9)

1. Moskovskiy filial Vsesoyuznogo instituta po proyektirova-  
niyu organizatsiy energeticheskogo stroitel'stva (for  
Vinit'skiy, Eberlin).

(Hoisting machinery)

(Electric power plants--Design and construction)



BYKOV, V.A.; GEL'FENBEYN, Ye.Yu.; PILIP, M.M.

New design of rope transfer and racking arrangements.  
Prokat. proizv. no.2:111-117 '60. (MIRA 14:11)  
(Rolling mills--Equipment and supplies)

DUNAYEV, V.I.; GEL'FENBEYN, Ye.Yu.

Removal of the finished product from plate mills. Prokat.  
proizv. no.2:118-125 '60. (MIRA 14:11)  
(Rolling mills—Equipment and supplies)

GRABOVSKIY, L.K., inzh.; BASHILOV, G.N., inzh.; SOKOLOVSKIY, O.P., inzh.;  
KRASNOSEL'SKIKH, S.N., inzh.; ANTONOV, P.A.; BYKOV, V.A., inzh.;  
DANILOV, G.G., inzh.; GEL'FENBEYN, Ye.Yu., inzh.; PILIP, M.M.,  
inzh.; MAKAROV, B.V., inzh.; RAGINSKIY, D.M., inzh.

Equipment of a working line of hot rolling mills. Sbor. st.  
NIITIAZHMASHa Uralmashzavoda no.6:70-96 '65.

(MIRA 18:11)

GEL'FENBOYM, Moisey Shlemovich, KOLCHANSKAYA, N.A., red.;  
MAMONTOVA, N.N., tekhn. red.

[Guide for the buyer of photographic supplies and photographic chemicals] Pokupateliu o fotomaterialakh i foto-khimikatakh. Moskva, Gostrogizdat, 1961. 81 p.

(MIRA 15:7)

(Photography--Equipment and supplies)

VARFOLOMEYEV, F.G.; GEL'FENBOYM, M.Sh.; KOTOVICH, Yu.V.;  
OSTANOVSKIY, T.S.; SEMENETS, V.M.; SHIROKOVA, Ye.A.;  
EYGINSON, Ye.N.; VVEDENSKIY, S.F., red.; SHEL'NIKOVA,  
TS.B., red.; TSESARKIN, L.D., red.

[Study of goods serving cultural needs] Tovarovedenie  
kul'ttvarov. [By] F.G.Varfolomeev i dr. Moskva, Izd-vo  
Ekonomika, 1964. 471 p. (MIRA 17:5)

GEL'FENSHTEYN, A.B.

Evaluation of the performance of the modern beer rectification  
columns with continuous action. Spirt.prom. 29 no.4:36-38 '63.  
(MIRA 16:5)

1. Karabaltinskiy spirtovoy zavod.  
(Distilling apparatus)

GEL'FENSHTEYN, A.B.; YERMOLENKO, R.T.

Karabalty Distillery is an enterprise of communist labor. Ferm.  
1 spirt.prom. 30 no.8:1-3 '64. (MIRA 18:1)

ORL'FER, A.P.

U-shaped pancreatocystojejunostomy in pancreatic cyst caused by  
injury. Khirurgiia no.10:88 0 '55. (MLRA 9:2)

1. Iz khirurgicheskogo otdeleniya 3-y gorodskoy bol'nitsy Kiyeva.  
(PANCREAS--TUMORS) (CYSTS)



GEL'FER, A.P., starshiy ordinator (Kiyev)

Phlegmon of the gastrointestinal tract. Vrach.delo no.5:505-509  
My '57. (MLRA 10:8)

1. Khirurgicheskoye otdeleniye (nachal'nik - kandidat meditsinskikh  
nauk D.N.Dumbadze) Bol'nitsy No.2 Yugo-zapadnoy zheleznoy dorogi  
(ALIMENTARY CANAL--DISEASES)

L 27581-66 ENT(1)/T RO/JK

ACC NR: AP6018376

SOURCE CODE: UR/0219/65/059/005/0087/0071

AUTHOR: Gel'fer, A. P.

ORG: Surgery Section /headed by Candidate of medical sciences D. N. Dugbadze/  
Railroad Hospital No. 2, South-Western Railroad (Khirurgicheskoye otdeleniye Dorozhnykh  
bol'nitsy No. 2 Yugo-zapadnoy zheleznoy dorogi); Toxicology Laboratory /headed by  
Doctor of medical sciences Yu. S. Kozan/, Kiev Institute of Labor Hygiene and Occupa-  
tional Diseases (Toksikologicheskaya laboratoriya Kiyevskogo instituta gigiyeny truda  
i profzabolevaniy)

TITLE: Experimental data on the effect of a number of organophosphorus compounds on  
the motility of the rabbit intestine

SOURCE: Byulleten' eksperimental'noy biologii i meditsiny, v. 59, no. 5, 1965, 67-71

TOPIC TAGS: organic phosphorus compound, rabbit, digestive system, pharmacology,  
muscle physiology

ABSTRACT: The effect of organophosphorus compounds on the motility of the  
rabbit intestine was investigated in situ upon intramuscular or intravenous  
injection of the compounds. The action of compounds of the following three  
groups was investigated: 1) those with a cholinesterase inhibitor action  
both in vitro and in vivo, i.e., pyrophos (I), isosystox (II), and methylace-  
tophos (III); 2) compounds which exhibit a relatively weak cholinesterase  
inhibitor effect in vitro, but are transformed into substances with a strong

Card 1/2

L 27581-66

ACC NR: AP6018376

anticholinesterase activity in vivo - i.e., octamethyl (IV), chlorophos (V), M-81 (VI), and M-81 Su 3 (VII); 3) K-20-35 (VIII), a substance which does not exert a cholinesterase inhibitor effect either in vitro or in vivo. Acetophos (IX) was also investigated. The action of the compounds on the motility of the intestine, expressed in ratios of increase of the amplitude of contractions of longitudinal muscle fibers to that of contractions of transverse muscle fibers, was as follows: I, 3:1.6; II, 2.5:3; III, 2.5:2.7; IV, 4.8:1.3; V, 5.1:4.8; VI, 4.5:2.5; VII, 4.6:2.2; VIII, 0:0; IX, 2:2.3. The increase in peristalsis was greatest when the induced contraction of longitudinal muscles exceeded that of transverse muscles to the greatest extent. Derivatives of pyrophosphoric acid (I, IV), esters of dithiophosphoric acid (VI, VII), and V (an ester of phosphinic acid) acted predominantly on longitudinal muscles, while esters of thiophosphoric acid (II, III, IX) had a greater effect on transverse vs. longitudinal muscles. III, VI, and VII increased the motility of the intestine in doses in which they produced only a weak anticholinesterase effect. Injection of III or IX after the motility of the intestine had been inhibited by a cholinolytic (atropine or tropacine) induced new contraction waves. This paper was presented by Active Member, AMN SSSR, A. I. Cherkas. Orig. art. has: 1 figure. [JPRS]

SUB CODE: 06 / SUBM DATE: 18Dec63 / ORIG REF: 010 / OTH REF: 002

Card 2/2

ACC NR: AF602903

SOURCE CODE: UR/0413/66/000/014/0024/0024

INVENTOR: Sanin, P. I.; Shepeleva, Ye. S.; Borodach, M. S.; Myannik, A. O.;  
Kagan, Yu. S.; Gel'fer, A. P.; Paykin, D. H.; Gamper, N. M.

ORG: none

TITLE: Preparation of esters of phosphoric and thiophosphoric acids. Class 12,  
No. 183751 [announced by Institute of Petrochemical Synthesis, AN SSSR (Institut  
neftekhimicheskogo sinteza AN SSSR)]

SOURCE: Izobret prom obraz tov zn, no. 14, 1966, 24

TOPIC TAGS: insecticide, chloroalkyl phosphate, chloroalkyl thiophosphate, ester,  
phosphoric acid

ABSTRACT: In the proposed method for the preparation of herbicides, the phos-  
phoric and thiophosphoric esters of the general formula:



(where X and Y are O or S; n = 1, 4, 6, 8; and R is an alkyl) are  
obtained by the reaction of trichloroalkyl alcohols with tetrachloro-  
alkanes [sic]. [WA-50; CBE No. 11]

SUB CODE: 07/ SUBM DATE: 21Jun65/

Card 1/1

UDC: 547.26'118.07

GEL'FER, A.P. [Hel'fer, A.P.]

Effect of some phosphorus organic compounds on the motor activity  
of the intestine. Fiziol. zhur. [Ukr.] 7 no.2:243-250 Apr '61.  
(MIRA 14:4)

1. Toxicological Laboratory of the Kiev Institute of Labor Hygiene  
and Occupational Diseases and the Surgical Division of Southwest  
Railway Hospital No.2.  
(INTESTINE MOTILITY) (PHOSPHORUS ORGANIC COMPOUNDS)

GEL'FER, A.P.

Experimental data on the effect of a series of organophosphorus compounds on the motor function of the intestines in rabbits. Biul.eksp.biol. i med. 59 no.5:67-71 '65.

(MIRA 18:11)

1. Khirurgicheskoye otdeleniye (nachal'nik - kand.med.nauk D.N.Dumbadze) Dorozhnoy bol'nitsy No.2 Yugo-zapadnoy zheleznoy dorogi i toksikologicheskaya laboratoriya (zav. - dok'tor med. nauk Yu.S.Kagan) Kiyevskogo instituta gigiyeny truda i professional'nykh zabolevaniy. Submitted December 18, 1963.

SASS, A.Ya., ~~CHILIKER~~ B.Ya., redaktor; YUZHNAIA, Ye.A., redaktor  
izdatel'stva; SOSNIN, A.P., tekhnicheskii redaktor

[SMD-2 peat-spreading machine] Stilochnaia mashina SMD-2.  
Moskva, Gos. izd-vo mestnoi promyshl. RSFSR, 1956.

135 p.

(MLRA 10:4)

(Peat machinery)

88428

S/056/60/039/006/015/063  
B006/B056

24.6900

AUTHORS: Vovchenko, V. G., Gel'fer, G., Kuznetsov, A. S.,  
Meshcheryakov, M. G., Svyatkovskiy, V.

TITLE: Effect of Nuclear Binding of Nucleons Upon the Shape of  
Pion Energy Spectra

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,  
Vol. 39, No. 6(12), pp. 1557-1570

TEXT: A description is given of experiments which were carried out with the aim of obtaining quantitative data on the effect produced by nucleon bindings in deuterons and carbon nuclei upon the production of charged pions. Conclusions are drawn with respect to pion production processes on the basis of comparisons of the energy spectra of pions produced in collisions of protons with free protons and with nucleons bound in deuterons and carbon nuclei. The experiments were conducted in a way ensuring strictly equal conditions in taking the spectra and separating the pp- and pn-collisions. The experiments were carried out on the six-meter synchrocyclotron of the Joint Institute of Nuclear

Card 1/6



88428

Effect of Nuclear Binding of Nucleons  
Upon the Shape of Pion Energy Spectra

S/056/60/039/006/015/063  
B006/B056

Research. For the pp- and pd-collision experiments, thin-walled cylindrical vessels (4 cm x 12 cm) filled with liquid hydrogen and deuterium, respectively, were used as targets. The densities were 0.0708 and 0.169 g/cm<sup>3</sup>, respectively. In the experiments with carbon a 3 mm thick, 5 x 5 cm<sup>2</sup> graphite plate was used as target. In the target center, the proton energy in all cases was (654±5) Mev, the slowing-down losses in hydrogen and deuterium, respectively, were 1.4 and 1.7 Mev, and in carbon they were 1.9 Mev; the slowing-down losses of the 150-Mev pions were 0.7, 0.8, and 0.5 Mev, respectively. The proton flux in a beam of 2 x 3 cm<sup>2</sup> cross section was 2·10<sup>8</sup> p/cm<sup>2</sup>·sec. The energy spectra of the charged pions were measured by a magnetic spectrometer with two thin scintillation counters at the input, which was described by L. S. Azhgirey et al. The pion recording threshold was about 35 Mev. The results obtained had, after a number of corrections, which had an error of 7% for the differential cross sections. The differential elastic pp scattering cross sections at 654 Mev and for 56° in the laboratory system (1.s.) were taken to be (6.7±0.35)·10<sup>-27</sup> cm<sup>2</sup>/steradian, so that for 120° in c.m.s. the value (3.41±0.13)·10<sup>-27</sup> cm<sup>2</sup>/steradian was obtained. The

Card 2/6

88428

Effect of Nuclear Binding of Nucleons  
Upon the Shape of Pion Energy Spectra

S/056/60/039/006/015/063  
B006/B056

difference found to exist in the spectra of mesons from deuterium and carbon is said to be due to the following reasons: a) a higher degree of pair correlation of the nucleons in the carbon nuclei than in deuterons, b) differences in nucleon momentum distribution in these nuclei, and c) effect of secondary pion-nucleon interactions in carbon nuclei. For an angle of  $\sim 90^\circ$  in the c.m.s. of the two colliding nucleons, the following ratio of differential cross sections was found:

$$\frac{d\sigma}{d\omega} [p + p \rightarrow \pi^+]_H : \frac{d\sigma}{d\omega} [p + p \rightarrow \pi^+]_D : \frac{d\sigma}{d\omega} [p + p \rightarrow \pi^+]_C = 1:0.79:0.40.$$

The  $\pi^-$ -meson yields from deuterium and carbon per nucleon of the target nucleus were found to be approximately the same. The  $\pi^+-\pi^-$ -meson yield ratios for deuterium and carbon were found to be  $10.3 \pm 1.3$  and  $6.0 \pm 0.8$ . The decrease of this ratio on the transition from deuterium to carbon is explained by the considerable fraction of secondary exchange interaction ( $\pi^0 + n \rightarrow \pi^- + p$ ) in the  $\pi^-$ -meson yield of carbon. B. S. Neganov, O. V. Savchenko, A. G. Meshkovskiy, Yu. D. Prokoshkin, L. B. Parfenov, and M. S. Kozodayev are mentioned. There are 4 figures,

Card 3/6

88428

Effect of Nuclear Binding of Nucleons  
Upon the Shape of Pion Energy Spectra

S/056/60/039/006/015/063  
B006/B056

3 tables, and 31 references: 14 Soviet, 14 US, 1 CERN, and 2 British.

ASSOCIATION: Ob'yedinenyy institut yadernykh issledovaniy  
(Joint Institute of Nuclear Research)

SUBMITTED: July 9, 1960

Text to Fig. 4 : Comparison between the energy spectra of the charged pions 1)  $\pi^+$ -mesons, emitted in free pp collisions. 2)  $\pi^+$  from  $[pp]_D$ ; 3)  $\pi^+$  from  $[pp]_C$ ; 4)  $\pi^-$  from  $[pn]_D$ ; 5)  $\pi^-$  from  $[pn]_C$ . All spectra are normalized on a uniform area.

Text to Table 1: Differential cross sections for charged pions in pp-pd- and pC-collisions at 654 Mev. Angle of observation  $56^\circ$  with respect to the proton beam ( $\sim 90^\circ$  in the c.m.s.). 1) Differential cross sections, in  $10^{-28}$  cm<sup>2</sup>/steradian; 2) Process; 3) Nucleons; 4) l.s.; 5) c.m.s.

Card 4/6

88428

S/056/60/039/006/015/063  
B006/B056




Fig. 4

X

Card 5/6

88428

S/056/60/039/006/015/063  
B 006/B056

Процесс 2	1 Дифференциальные сечения, $10^{-38}$ см <sup>2</sup> /стерад			
	3 $\frac{d\sigma^+}{d\Omega}, 56^\circ$ (я.с.)	4 $\frac{d\sigma^-}{d\Omega}, 56^\circ$ (я.с.)	5 $\frac{d\sigma^+}{d\Omega}, \sim 60^\circ$ (с.п.м.)	5 $\frac{d\sigma^-}{d\Omega}, \sim 60^\circ$ (с.п.м.)
$p+p \rightarrow \pi^+ + \text{нуклоны}$ 3	$10,2 \pm 1,0$	—	$6,7 \pm 0,7$	—
$p+d \rightarrow \pi^+ + \text{нуклоны}$ 3	$9,1 \pm 0,9$	$0,88 \pm 0,12$	$5,9 \pm 0,6$	$0,57 \pm 0,08$
$p+C \rightarrow \pi^+ + \text{нуклоны}$ 3	$30,2 \pm 3,0$	$5,0 \pm 0,8$	$19,5 \pm 1,9$	$3,2 \pm 0,5$

Tab. 1

Card 6/6

GEL'FER, G.A.

Essential hypotension; its distribution and clinical course. Ter. arkh.,  
Moskva 24 no.4:53-60 July-Aug 1952. (CINL 23:2)

1. Of the Department of Propedeutics of Internal Diseases (Director --  
Prof. K. G. Nikulin), Gor'kiy Medical Institute imeni S. M. Kirov.

GEL'FER, G.A., dotsent

Remote results of treating hypertension with prolonged sleep. Terap.  
arkh. 26 no.5:29-33 8-0 '54. (MLRA 8:2)

1. Iz propedevticheskoy terapevticheskoy kliniki (dir. prof. K.O.  
Nikulin) Gor'kovskogo meditsinskogo instituta imeni S.M.Kirova  
(SLEEP, therapeutic use,  
hypertension, results)  
(HYPERTENSION, therapy,  
sleep, results)

GEL'FER, G.A. (Gor'kiy)

Unconditioned vascular reactions in hypertension and their dynamics  
in sleep therapy. Klin. med. 32 no.6:60-66 Je '54. (MLRA 7:8)

1. Iz kafedry propedevtiki vnutrennikh bolezney (zav.-prof. K.G.  
Nikulina) Gor'rovskogo meditsinskogo instituta.

(HYPERTENSION, therapy,

\*sleep ther., eff. on unconditioned vasc. reactions)

(BLOODVESSELS

\*hypertension, unconditioned reflexes, eff. of sleep ther.)

(SLEEP, therapeutic use

\*hypertension, eff. on unconditioned reflexes of blood  
vessels)



PHASE I BOOK EXPLOITATION

SOV/4948

Gel'fer, Gesel' Ayzikovich, Aleksandr Vladimirovich Ivanov, and Yakov  
Grigor'yevich Medvedev

Vzryvozashchishchennoye elektrooborudovaniye: spravochik dlya rabotnikov  
neftepererabatyvayushchey i gazovoy promyshlennosti (Explosionproof  
Electrical Equipment: Manual for Oil-Refinery and Gas-Industry Workers)  
Leningrad, Gostoptekhizdat, 1960. 328 p. Errata slip inserted. 4,100  
copies printed.

Ed.: V. Ye. Ul'yashchenko; Tech. Ed.: P. S. Frumkin; Executive Ed.:  
P. S. Dolmatov.

**PURPOSE:** This manual is intended for engineers and technicians working in  
oil refineries and in the gas industry, and may be useful to personnel  
in other industries where the hazard of gas or dust explosion exists.

**COVERAGE:** The manual contains the specification and description of ex-  
plosionproof electric machines, apparatus, and devices manufactured by  
Soviet industry. Data on classification of locations by the degree of

Card 1/13

Explosionproof Electrical Equipment (Cont.)

SOV/4948

explosion hazard, classification of explosive mixtures, selection and use of explosionproof equipment, and arrangement of electric networks in locations containing explosive substances are discussed. It is stated in the foreword that this manual is the first attempt to present in a systematic way data relating to explosionproof electrical equipment and its use. It is based on the Soviet "Rules for the Arrangement of Electrical Installations," 1957-1958 edition, the directives and instructions of the former Ministry of Petroleum and Chemical Industries, USSR, and various scientific research and planning institutes, and information from manufacturing plants. The manual uses the terms *vzryvozashchishchennyy* (protected against explosion), *vzryvobezopasnyy* (explosion-safe) and *vzryvonepronitsayemyy* (impenetrable to explosion) but does not make clear what difference in meaning, if any, exists. All three terms are hereafter translated as "explosionproof." No personalities are mentioned. There are 22 references, all Soviet.

Card ~~2/13~~

GEL'FER, G.A., dotsent

Diagnostic value of electrophoretic determination of blood  
protein fractions in stenocardia. Terap.arkh. 32 no.12:42-47  
'60. (MIRA 14:2)

1. Is kafedry propedevtiki vnutrennikh bolezney (zav. - prof.  
K.G. Nikulin) Gor'kovskogo meditsinskogo instituta imeni  
S.M. Kirova.  
(BLOOD PROTEINS) (ANGINA PECTORIS)

BOLDIN, K.M. (Yaroslavl'); DROZDOVA, Z.S.; LEVIN, R.I.; VAYSMAN, L.A. (Kuybyshev-obl.); PODOSINOVSKIY, V.V. (Kazan'); SAYFULLINA, Kh.M. (Kazan'); EUSYGIN, N.V. (Kazan'); RAZUMOVSKIY, Yu.K. (Leninogorsk); GEL'FER, G.A., dotsent (Gor'kiy); MAMISH, M.G. (Kazan'); RAFALOVICH, M.B., dotsent; MEL'NICHUK, S.P., kand.med'nauk; KRAPIVIN, B.V.; STAROVEROV, A.T. (Saratov); SURIN, V.M.; PORosenkov, V.S. (Romodanovo, Mordovskoy ASSR); ANDROSOV, M.D. (Moskva); ZARIPOV, Z.A. (Urussu, Tatarskoy ASSR); MURAV'YEV, M.F. (Izhevsk); KUZ'MIN, V.I. (Batyrevo, Chuvashskoy ASSR); SITDYKOV, E.N. (Kazan'); YUDIN, Ya.B. (Novokuznetsk)

Short reports. Kaz.med.zhur. no.4:81-91 J1-Ag '62. (MIRA 15:8)  
(MEDICINE--ABSTRACTS)

KSENZ, S.P.; GRUNICHEV, A.S., kand. tekhn. nauk, retsenzent;  
KRASNOV, I.F., kand. tekhn. nauk, retsenzent; GEL'FER, I.N.,  
red.; KOCHETKOVA, N.A., red.

[Searching for faults in radioelectronic systems using a  
functional test method] Poisk neispravnostei v radioelektron-  
nykh sistemakh metodom funktsional'nykh prob. Moskva, Sovet-  
skoe radio, 1965. 135 p. (MIRA 18:4)

AUTHOR: Gel'fer, L.M., Engineer SOV/117-58-11-8/36

TITLE: ~~An Accelerated~~ Cross Motion on a Turret Lathe (Uskorennaya poperechnaya podacha na revol'vernom stanke).

PERIODICAL: Mashinostroitel', 1958, Nr 11, p 10 (USSR)

ABSTRACT: In the Khar'kovskiy elektromekhanicheskiy zavod imeni Stalina (Khar'kov Electromechanical Plant imeni Stalin), the turret lathe model 136 has been modernized by the installation of a special device with a separate motor of 0.36 kw and 1,440 rpm. This device accelerates the cross motion of the lathe and cuts the time needed for auxiliary operation. There is 1 diagram.

1. Lathes--Operation 2. Lathes--Equipment

Card 1/1

TAYTS, Ye.M., doktor tekhn. nauk; SHVARTS, S.A., kand. tekhn. nauk[deceased]; PEYSAKHZON, I.B., inzh.; GEL'FER, M.L., inzh.; DMITRIYENKO, M.T., inzh.; DORFMAN, G.A., inzh.; IZRAELIT, Ye.M., inzh.; KULAKOV, N.K., inzh.; KUSHLYANSKIY, B.S., inzh.; MEYKSON, L.V., inzh.[deceased]; LEONOV, A.S., inzh.; SHVARTS, G.A., inzh.; SHVARTSMAN, I.Ya., inzh.; YATSENKO, N.Ya., inzh.; BABIN, P.P., inzh.; KHANIN, I.M., doktor tekhn. nauk, prof., red.; KOZYREV, V.P., inzh., red., KUPENMAN, P.I., inzh., red.; LGALOV, K.I., inzh., red.; LEYTES, V.A., inzh., red.; LERNER, B.Z., inzh., red.; POTAPOV, A.G., inzh., red.; SHELKOV, A.K., red.

[By-product coke industry worker's handbook in six volumes]  
Spravochnik koksokhimika v shesti tomakh. Moskva, Metal..  
lurgiya. Vol.2. 1965. 288 p. (MIRA 18:8)

MARTYNYUK, Arseniy Grigor'yevich, professor; GEL'FER, P.I., redaktor;  
LOKHMATYY, Ye.G., tekhnicheskiy redaktor

[Form and function of the kidney surviving unilateral nephrectomy]  
Sostoianie i funktsia ostavsheisia pochki. Kiev, Gos.med. izd-vo  
USSR, 1955. 87 p. (MLA 9:3)

(KIDNEYS)



GEL'FER, P.I., professor (Kiyev)

"Outlines of operative urology" by A.P.TSulukidze. Reviewed by  
P.I.Gel'fer, Urologia 21 no.2:84-86 Ap-Je '56. (MIRA 9:12)  
(OMNITOURINARY ORGANS--SURGERY)  
(TSULUKIDZE, A.P.)

GEL'FER, P.I., prof.; BLATNOY, Kh.P.

Modification of single-stage prostate adenomectomy with primary  
suture of the urinary bladder. Urologiya 24 no.4:33-36 J1-Ag '59.

(MIRA 12:12)

1. Iz Kiyevskoy oblastnoy bol'nitsy (glavnyy vrach A.M. Cherednik).  
(PROSTATIC HYPERTROPHY surgery)  
(BLADDER surgery)

KUZMENKO, Leonid Nikolayevich, prof.; GEL'FER, P.I., red.; OITSHTAYN,  
A.D., tekhnred.

[Urolithiasis] Mochekamennaya bolezni'. Kiev, Gos.med.izd-vo  
USSR, 1960. 287 p. (MIRA 13:12)  
(CALCULI, URINARY)

VEYMEROV, Isaak Borisovich, prof.; ROZHINSKIY, Lazar' Markovich;  
GEL'FER, P.I., red.; GITSHEYN, A.D., tekhn. red.; CHUCHUPAK,  
V.D., tekhn. red.

[Diseases of the male urogenital organs; the lower section of  
the urogenital system] Bolezni mocheopolovykh organov u muzh-  
chin; nizhnii otdel mocheopolovoi sistemy. Kiev, Gos. med.  
izd-vo USSR, 1961. 220 p. (MIRA 15:3)  
(GENITOURINARY ORGANS—DISEASES)

GEL'FER, P.I.

Hematuria in prostatic adenoma. Urologia 26 no.2:23-27 '61.  
(PROSTATE GLAND--TUMORS) (MIRA 14:3)  
(HEMATURIA)

GEL'FER, P.I.

"Pyelonephritis" by A.IA. Pytel', S.D. Goligorskii. Reviewed  
by P.I. Gel'fer. Urologiia no.1:90-92 '62. (MIRA 15:11)  
(KIDNEYS—DISEASES) (PYTEL', A.IA.) (GOLIGORSKII, S.D.)

GEL'FER, P.I., prof. (Kiyev)

Current drugs used in urology. Urologiia 27 no.4:66-70 J1-Ag  
'62. (MIRA 15:11)

(UROLOGY)

(DRUGS)

POLONSKIY, B.L., prof., red.; PROSKURA, O.V., dots., red.; ALAPIN, G.Ya., prof., red.; GEL'FER, P.I. (Kiev), red.; PINEVICH, M.V., dots., doktor med. nauk (Vinnitsa); TSYBUL'SKIY, L.Ye., red.; MARINSKAYA, A.L., tekhn. red.

[Transactions of the Ukrainian Conference of Urologists devoted to the 150th anniversary of N.I.Pirogov's birth, held June 27-29, 1960] Trudy Ukrainskoi respublikanskoi konferentsii urologov, posviashchena 150-letiiu so dnia rozhdeniia N.I. Pirogova, 1960. Kiev, Gosmedizdat USSR, 1962. 386 p.

(MIRA 16:3)

1. Ukrainskaya respublikanskaya konferentsiya urologov, posvyashchena 150-letiyu so dnya rozhdeniya N.I.Pirogova, 1960.
2. Glavnyy urolog Ministerstva zdavookhraneniya Ukr.SSR (for Proskura).

(UROLOGY--CONGRESSES)



ALAPIN, G.Ya., prof., red., (Khar'kov); GEL'FER, F.I., prof.,  
red.; FINEVICH, M.V., dots., red.; POLONSKIY, B.L., prof.,  
red.; PROSKURA, O.V., dots., red.; TSYBUL'SKIY, L.Ye.,  
red.; NARINSKAYA, A.L., tekhn. red.

[Transactions of the Republic Conference of Urologists  
(dedicated to the 150th anniversary of N.I.Pirogov's birth)]  
Trudy Respublikanskoi konferentsii urologov (posviashchena  
150-letiiu so dnia rozhdeniia N.I.Pirogova) 27-29 iiunia 1960.  
Gosmedizdat, USSR, 1962. 386 p. (MIRA 16:12)

1. Respublikanskaya konferentsiya urologov Ukrainskoy SSR,  
1960.

(UROLOGY)

GEL'FER, P.I., prof.

Review of B.S.Gekhman's book "Nonspecific epididymitis." Vest.  
derm. i ven. 38 no.9:82-83 S '64. (MIRA 18:4)

GEL'FER, P.I., prof. (Kiyev)

Current aspects of the treatment of urolithiasis;  
the so-called dissolution of stones. Urol. i nefr.  
no.2:64-71 '65. (MIPA 19:1)

RUETSOVA, L. K., GANDURO, R. L., EYDEL'SHTEYN, S. I., GEL'FER, H. A.

Penicillin inhalation therapy. *Pediatrics*, Moskva No. 6,  
Nov.-Dec. 50. p. 44-51.

1. Of the Department of Pediatrics, Central Institute for the  
Advanced Training of Physicians (Head of Department--Prof.  
G. N. Speranskiy, Active Member of the Academy of Medical Sciences)  
attached to the Hospital imeni Dzorzhinskiy (Head Physician--  
Ye. L. Guterman) and of the Department of Experimental Therapy  
VNIP (Head of Department--Prof. Z. V. Yermol'yeva, Corresponding  
Member of the Academy of Medical Sciences).

CLM. 20, 3, March 1951

ZHURAKHOVSKIY, P., inzh.; GEL'FER, S., inzh.; GAYETSKIY, A., inzh.

Machine for cutting adhesive rubber. Avt.transp. 40 no.10:31  
0 '62. (MIRA 15:11)

(Cutting machines)

ZHURAKHOVSKIY, P.; GEL'FER, S.; GAYETSKIY, A.

Machine for inserting vulcanization devices. Avt.transp. 40  
no.4:53 Ap '62. (MIRA 15:4)  
(Vulcanization--Equipment and supplies)

ZHURAKHOVSKIY, P.N.; GEL'FER, S.; GAYETSKIY, A.

Glue spraying unit. Avt.transp. 4 no.8:52-53 Ag '62. (MIRA 16:4)  
(Gluing--Equipment and supplies)

GEL'FER, S.A.

Variation method in conformal mapping. Izv. AN Arm. SSR. Ser.  
fiz.-mat. nauk 14 no.3:17-24 '61. (MIRA 14:8)

1. Gor'kovskiy inzhenerno-stroitel'nyy institut imeni V.P.  
Chkalova.

(Conformal mapping)



ZHURAKHOVSKIY, R.; GAYETSKIY, A.; GEL'FER S.

Tire-retreading line. Avt.transp. 41 no.4135-36 Ap '63.  
(MIRA 16:5)  
(Tires, Rubber--Retreading and recapping)

GELFER, S.A.

Gelfer, S. On the class of regular functions which do not take on any pair of values  $\alpha$  and  $\beta$ . *Russ. Math. Math. Sbornik* [N.S. 19(61), 33-5] (1946). (Russian; English summary)

The author considers the following two classes of functions: the class (C) of functions  $w = f(z) = \sum_{n=0}^{\infty} a_n z^n$  which are regular in  $|z| < 1$  and satisfy the condition that  $f(z_1) \cdot f(z_2) \neq 1$  for  $|z_1| < 1, |z_2| < 1$ , the class (D) of functions  $w = \varphi(z) = 1 + \sum_{n=1}^{\infty} a_n z^n$  which are regular in  $|z| < 1$  and satisfy  $\varphi(z_1) + \varphi(z_2) \neq 0$  for  $|z_1| < 1, |z_2| < 1$ . If  $f$  belongs to (C), then  $(1+f(z))/(1-f(z))$  belongs to (D); conversely, if  $\varphi$  belongs to (D),  $(\varphi(z)-1)/(\varphi(z)+1)$  belongs to (C). The author finds inequalities for the moduli of these functions and their derivatives and on the basis of these he obtains results to obtain known properties of these functions.

*H. C. Spencer* (see also 1946d, 1946e, 1946f)

*Mathematical Reviews,*

Vol

No.

Gelf'fer, S. A.

USSR.

1-P/W

Gelf'fer, S. A. On typically real functions of order  $p$ .  
Mat. Sb. N.S. 35(77), 193-214 (1954). (Russian)

Let  $T(p)$  denote the class of functions  $\sum_{k=0}^{\infty} a_k z^k$  which are regular in  $|z| < 1$ , have all coefficients real, and for which there is a  $\rho < 1$  such that  $\Im f(z)$  changes sign  $2p$  times on each circle  $|z| = r$ ,  $\rho < r < 1$ . The reviewer and Robertson [Trans. Amer. Math. Soc. 70, 127-136 (1951); MR 12, 691] proved that if  $f(z) \in T(p)$  then for each  $n > p$

$$(1) \quad |a_n| \leq \sum_{k=1}^p |a_k| \frac{2k(n+p)!}{(n-k)!(p+k)!(p-k)!(n-p-1)!}$$

and this inequality is sharp in all of the variables.

The author now gives a second proof of (1) which is somewhat longer than the original. It is proved that if  $f(z) \in T(p)$ , then it can be represented in the form

62

1/2

(over)

*Goodman, S. A.*

$$(2) \quad f(s) = \sum_{n=1}^{\infty} A^{(n)} \prod_{k=1}^n s(\cos \theta_k) + \frac{1}{\pi} \prod_{k=1}^n s(\cos \theta_k) \int_0^\pi s(\cos \theta) d\mu(\theta),$$

where  $A^{(n)}, \dots, A^{(n-1)}$  are real,  $\mu(\theta)$  is a real non-decreasing function and  $s(x) = x/(1-2x^2+x^4)$ . The class of functions given by the right side of (2) is wider than  $T(p)$ , so the bound (1) is now extended to this wider class.

A. W. Goodman (Lexington, Ky.).

Gel'fer, S. A. On coefficients of typically real functions.

Doklady Akad. Nauk SSSR (N.S.) 94, 373-376 (1954).

(Russian)

Let  $F(z) = \sum_{n=1}^{\infty} a_n z^n$  be typically-real in  $0 < |z| < 1$ . It is proved that (a) if  $a_1 - a_{-1} = 1$  then  $|a_n| \leq n$ ,  $n \geq 2$ , and (b) if instead  $a_{-1} = -1$  and  $F(z) \neq 0$  in  $|z| < 1$  then  $-2 \leq a_n \leq 2$ ,  $-1 \leq a_1 \leq 3$ , and

$$4 \min_{0 \leq \theta \leq 2\pi} \sin n\theta \sin \theta \leq a_n \leq 4 \max_{0 \leq \theta \leq 2\pi} \sin n\theta \sin \theta \quad \text{for } n \geq 2.$$

All the inequalities are sharp. In the proof of (b) the author uses a Stieltjes integral representation which he attributes to Goluzin, but which was proved earlier by M. S. Robertson [Bull. Amer. Math. Soc. 41, 565-572 (1935)]. The result (a) is partly contained in a more general theorem of Nehari and Schwarz [see the preceding review].

A. W. Goodman (Lexington, Ky.).

62

GELFER, S.H.

USSR/Mathematics - Multi-leafed functions

Card 1/1 : Pub. 22 - 1/44

Authors : Golfer, S. A.

Title : Variations of multi-leafed functions

Periodical : Dok. AN SSSR 98/6, 885-888, October 21, 1954

Abstract : A method for obtaining the formulas of a variation of multi-leafed functions is described. The method consists in reducing all operations to the determination of the variation of a function which maps a circle on single-leafed region of a plane (Riemannian) with uniforming variable. The theorem presented gives all the necessary definitions. Two references (1943 and 1946).

Institution : Gorkiy Engineering-Constructional Institute im V. P. Chkalov

Presented by: Academician M. A. Lavren'ev, July 19, 1954.

GEL'FER, S.A.

SUBJECT USSR/MATHEMATICS/Theory of functions CARD 1/1 PG - 548  
 AUTHOR GEL'FER S.A.  
 TITLE Variation method in the theory of the p-sheeted functions.  
 PERIODICAL Uspechi mat.Nauk 11, 5, 60-66 (1956)  
 reviewed 1/1957

Let  $F_p$  be the class of functions  $z = f(\zeta) = \sum_{n=1}^{\infty} c_n \zeta^n$ ,  $|\zeta| < 1$ , which map

the unit circle onto a region of the p-sheeted Riemannian surface.

Let  $F_p^g(a_1, \dots, a_m)$  be a function family of  $F_p$ . The functions belonging to it have the following properties: 1. they map the unit circle onto a region D of a Riemannian surface the genus of which is at most g. Here all functions which map the circle  $|\zeta| < 1$  onto regions of the same Riemannian surface, transfer the point  $\zeta = 0$  into the same point of the Riemannian surface which lies over  $z = 0$ . 2. In  $|\zeta| < 1$ ,  $f(\zeta)$  does not assume the values  $a_1, \dots, a_m$ .

The principal result of the paper is the following theorem: For given  $n \geq 1$  let  $|c_n|$  reach a maximum for the function  $\tilde{f}(\zeta) \in F_p^g(a_1, \dots, a_m)$ . Then  $\tilde{f}(\zeta)$  maps the unit circle  $|\zeta| < 1$  onto the whole Riemannian surface R which is cut along a finite number of analytic arcs. These intersections go through all points of  $R_1$  which lie over  $z = a_1, \dots, a_m, \infty$ .

INSTITUTION: Gorkij.

Call Nr: AF 1108825

Transactions of the Third All-union Mathematical Congress \* (Cont.) Moscow  
Jun-Jul '56, Trudy '56, V. 1., Sect. Rpts., Izdatel'stvo AN SSSR, Moscow, 1956, 237 pp.  
Gagayev, B. M. (Kazan'). Some Properties of Orthogonal Functions. 77

Gakhov, F. D. (Rostov-na-Donu); Krikunov, Yu. M. (Kazan').  
Topological Methods in the Theory of Function of a Complex  
Variable and Their Application for Solving Inverse  
Boundary Problems. 77-78

There is 1 reference, which is a translation into Russian.

Gel'fer, S. A. (Gor'kiy). On a Maximum Conformal Radius  
of the Fundamental Domain of a Given Group. 78

Mention is made of Lavrent'yev, M. A.

Geronimus, Ya. L. (Khar'kov). On Some Sufficient Convergence  
Conditions of the Fourier-Chebyshev Process. 78-79

Card 24/80

\*



Gelfer, S. A. On the coefficient problem for  $p$ -valent functions. Dokl Akad Nauk SSSR 196 955-958 (Russian)

Let  $F_p(a_1, \dots, a_n)$  denote the class of functions  $f(z) = \sum_{k=1}^{\infty} a_k z^k$  which are regular in  $|z| < 1$  and map onto a region  $D$  lying in some  $p$ -sheeted universal Riemann surface  $R$  of genus  $g$  and omit  $a_1, a_2, \dots, a_n$ ,  $n < 1$ . Using variational formulas developed earlier by the author [same Dokl 98 (1954), 645, 646; MR 16:417] he now proves that if  $f(z) \in F_p(a_1, \dots, a_n)$  and  $|f(z)| \leq 1$  on the entire surface  $R$  except for a finite number of analytic arcs. A. W. Goodman

*Goodman*

Gor'kovskiy Inzhenerno-stroitel'nyy institut imeni V. P. Chkalova.

GEL'FER, S. A.

20-2-2/62

AUTHOR: Gel'fer, S. A.

TITLE: On the Coefficients of the Typically Real Functions.  
(O koefitsiyentakh tipichno-veshchestvennykh funktsii)

PERIODICAL: Doklady Akad. Nauk SSSR, 1957, Vol. 115, Nr 2, pp. 211-213 (USSR)

ABSTRACT: A function  $f(z)$  is called typically real in the circle  $|z| < 1$ , when it is real in the case of a real  $z$  and when it satisfies in the other points of this circle the condition  $\operatorname{Im}(f(z)) \cdot \operatorname{Im}(z) > 0$ .  $T$  be a class of typically real functions  $f(z)$  which are regular in  $|z| < 1$  and satisfy the normalization conditions  $f(0)=0$ ,  $f'(0)=1$ . The present report examines the subclass  $T^{(2)}$  of the function  $f_2(z) \in T$ , for which  $f_2(z) = (1/i)f_2(iz) \in T$  applies. These functions satisfy the conditions  $\operatorname{Im}(f(z)) \cdot \operatorname{Im}(z) > 0$  and  $\operatorname{Re}(f_2(z)) \cdot \operatorname{Re}(z) > 0$  and all of them are odd. The author here puts down  $f_2(z) = z + x_1 z^3 + x_2 z^5 + \dots + x_n z^{2n+1} + \dots$ . This class  $T^{(2)}$  contains all odd one-sheet functions of the type  $f_2(z)$  with real coefficients. For the coefficients  $x_n$  the following exact estimations are known:  
 $|x_n| + |x_{n-1}| \leq 2 \quad (n=2,3,\dots); \quad -1 \leq x_1 \leq 1, -1/2 \leq x_2 \leq 3/2$ . There exist no exact estimations of the individual coefficients  $x_n$  for  $n > 2$ . Then the following theorem is given: In the case of  $f_2(z) \in T^{(2)}$  the following applies:

Card 1/2

GEL'FER, S. A.

20-2-2/60

AUTHOR: Gel'fer, S. A.

TITLE: On the Maximum of the Conformal Radius of the Fundamental Domain of a Double-Periodical Group (O maksimume konformnogo radiusa fundamental'noy oblasti dvoyako-periodicheskoy gruppy)

PERIODICAL: Doklady Akademii Nauk SSSR, 1957, Vol. 114, Nr 2, pp.241-244 (USSR)

ABSTRACT: Be it assumed that  $\{D\}$  is a family of simply connected domains  $D$  of the plane  $w$ , which contain the point  $w = 0$  and possess the following properties: 1) The domain  $D$  contains no points that are congruent with respect to the group  $T$  of the transformations  $w' = w + n_1 \omega_1 + n_2 \omega_2$ . (Here  $\omega_1$  and  $\omega_2$  denote constants with nonreal relations;  $n_1$  and  $n_2$  denote any whole numbers). The domain  $D$  does not contain the assumed system of the finite points  $a_1, \dots, a_m$  and the points congruent with respect to the group  $T$ . From all domains  $\{D\}$  that domain has to be determined here which has the highest conform radius. The class of the functions  $w = f(z) = \sum_{n=-\infty}^{\infty} c_n z^n$ , which are regular in the circle

Card 1/2

20-2-2/60

On the Maximum of the ~~Conformal~~ Radius of the Fundamental Domain of a Double-Periodical Group

$|z| < 1$  and represent this circle one-leaved onto the domains of the family  $\{D\}$ , are denoted here by  $S(w_1, w_2)$ . This problem is reduced here to the determination of the maximum  $|f'(0)| = |o_1|$  within the class  $S(\omega_1, \omega_2)$ . At first a rather voluminous theorem is given and proved. Also the theorems and conclusions resulting from this first theorem are given. There are 1 figure, and 5 Soviet references.

ASSOCIATION: Constructional Engineering Institute imeni V. P. Chkalov, Gor'kiy (Gor'kovskiy inshenerno-stroitel'nyy institut im. V. P. Chkalova)

PRESENTED: November 29, 1956, by V. I. Smirnov, Academician

SUBMITTED: September 23, 1956

AVAILABLE: Library of Congress

Card 2/2

GEL'FER. S. A. Doc Phys-Math Sci -- (diss) "Extreme problems in the theory of ~~multivariate~~ <sup>multivariate</sup> functions." Len, 1958. 17 pp (Len Order of Lenin State Univ im A. A. Zhdanov), 150 copies. Bibliography at end of text (12 titles).  
(KL, 11-58, 111)

-2-

AUTHOR: Gel'fer, S.A. (Gor'kiy)

39-44-2-4/10

TITLE: On the Maximum of the Conformal Radius of the Fundamental Domain of a Given Group (O maksimume konformnogo radiusa fundamental'noy oblasti dannoy gruppy)

PERIODICAL: Matematicheskii Sbornik, 1958, Vol 44, Nr 2, pp 213-223 (USSR)

ABSTRACT: Let  $S_a(\omega_1, \omega_2)$  be the class of the functions

$$w = f(z) = \sum_{n=1}^{\infty} c_n z^n,$$

which are regular in  $|z| < 1$  and which map this circle schlicht onto simply connected domains  $D$  which possess the following properties: Let  $T_n$  be the group of the transformations

$w' = w + n_1 \omega_1 + n_2 \omega_2$ , where  $\omega_1$  and  $\omega_2$  are constant,  $\frac{\omega_1}{\omega_2}$

not real, and  $n_1, n_2$  integer; then  $D$  contains no points which are congruent with regard to  $T_n$ , furthermore  $D$  does not contain the points  $a_1, \dots, a_m$  and those which are congruent to them with regard to  $T_n$ , and finally  $D$  contains the zero point.

Card 1/3

On the Maximum of the Conformal Radius of the Fundamental Domain of a Given Group

32-44-2-4/10

Theorem 1: If for  $w = f(z) \in S_n(\omega_1, \omega_2)$  the functional  $|f'(0)|$  attains a maximum, then  $w$  satisfies the differential equation

$$\frac{1}{z^2 w'^2} = A_0 + \sum_{i=1}^E A_i \zeta(w - a_i) + A_{m+1} \zeta(w) + \wp(w),$$

where  $A_i$  are constants,  $\sum_{i=1}^{m+1} A_i = 0$ , and  $\zeta$  and  $\wp$  are the Weier-

strasz  $\zeta$  and  $\wp$ -functions. Furthermore  $f(z)$  maps the unit circle onto a domain  $D$  with the following properties:

- 1.)  $D$  contains  $w = 0$ , it is the simply connected fundamental domain  $S_0$  of the group  $T_n$  with sections. The boundary of  $D$  consists of a finite number of analytic arcs which are mutually congruent with regard to  $T_n$ , and of piecewise analytic sections which connect the boundary with the points  $a_i$  ( $i=1, \dots, m$ ).
- 2.) To each pair of congruent curves of the boundary  $S_0$  and to the simple curves of the sections there correspond two curves of equal length on  $|z| = 1$ .

Card 2/ 3

On the Maximum of the Conformal Radius of the Fundamental Domain of a Given Group

39-44-2-4/10

Theorem 2: The domain  $D$  with the properties 1 and 2 of theorem 1 is uniquely determined.  
The theorems are needed for the determination of that domain  $D$  which possesses the greatest conformal radius, a problem which can be considered as a generalization of the well-known theorem of Koebe and of an investigation of Lavrent'yev [Ref 1] .  
Several special cases are considered. There are 2 figures and 4 Soviet references.

SUBMITTED: October 29, 1956

AVAILABLE: Library of Congress

1. Conformal mapping
2. Differential equations

Card 3/3



16(1)

AUTHOR: Gel'fer, S.A.

30V/20-126-3-1/69

TITLE: On the Maximum Conformal Radius of the Fundamental Region for the Group of Fractional Linear Transformations

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 126, Nr 3, pp 463-466 (USSR)

ABSTRACT: Let  $\{T\}$  be the proper discontinuous group of the fractional linear transformations  $T$ . Let  $\{D\}$  be a family of simply connected domains  $D$  of the  $w$ -plane with the following properties: 1)  $D$  contains no points congruent with respect to  $\{T\}$ ; 2)  $D$  does not contain the given point system  $a_1, \dots, a_m, \infty$  and the points congruent to this point with respect to  $\{T\}$ ; 3)  $D$  contains the point  $o$  which is different from the fixed points and the cyclic points of the group transformations. Let  $S_a(T)$  be the class of

functions  $w = f(z) = \sum_{n=0}^{\infty} c_n z^n$  which are regular in  $|z| < 1$  and

which map conformally the circle  $|z| < 1$  onto the domains  $\{D\}$ .

Theorem: If the functional  $|f'(0)|$  gets a maximum by the function  $f(z) \in S_a(T)$ , then the function maps the circle  $|z| < 1$

onto a domain  $D$  with the following properties: 1.  $D$  contains  $o$

and is a simply connected fundamental domain  $S_o$  of the group  $\{T\}$

Card 1/2

On the Maximum Conformal Radius of the Fundamental Region for the Group of Fractional Linear Transformations SOV/20-126-3-1/69

with cuts. The boundary of  $D$  consists of finitely many analytic arcs pairwise congruent with respect to  $\{T\}$ , and of piecewise analytic cuts beginning at the boundary of  $S_0$  and ending in the points  $a_1, \dots, a_m$  or in points congruent with them. 2. To every pair of congruent arcs of the boundary of  $S_0$  and to the simple arcs of the cuts on  $|z|=1$  for the mapping  $w = f(z)$  there correspond two arcs of the same length. 3. For the given group  $D$  is determined uniquely. There follow several examples for the application of the theorem for the determination of domains with a maximal conformal radius with respect to  $c_0$ .

The author mentions G.M. Goluzin, and M.A. Lavrent'yev. There are 6 Soviet references.

ASSOCIATION: Gor'kovskiy inzhenerno-stroitel'nyy institut imeni V.P. Chkalova (Gor'kiy Civil Engineering Institute imeni V.P. Chkalov)

PRESENTED: February 9, 1959, by V.I. Smirnov, Academician

SUBMITTED: February 2, 1959

Card 2/2

85944

16.3000

S/039/60/052/001/006/009 XX  
C111/C222

AUTHOR: Gel'fer, S.A. (Gor'kiy)

TITLE: On the Maximum of the Conformal Radius of the Fundamental Region  
of the Group of Fractional-Linear Transformations 16

PERIODICAL: Matematicheskii sbornik, 1960, Vol. 52, No. 1, pp. 629-640

TEXT: Let  $\{T\}$  be a properly discontinuous group of fractional-linear transformations and  $a_1, \dots, a_n$  be a system of points in the fundamental region of the group. Let  $\{D\}$  be a family of simply connected regions of the  $w$ -plane with the properties:  
1)  $D$  contains no points congruent with respect to  $\{T\}$ ; 2)  $D$  does not contain the points  $a_1, \dots, a_n, \infty$  and the points congruent to them with respect to  $\{T\}$ ; 3)  $D$  contains a fixed point  $c_0$  different from the fixed points and the cyclic points of the transformations of the group.  
Problem: Among all  $D$  of the family  $\{D\}$ , that region shall be determined which has the greatest conformal radius with respect to  $c_0$ .

Let  $S_a(T)$  be the class of functions

Card 1/4

85944

On the Maximum of the Conformal Radius of the  $S/039/60/052/001/006/009$  XX  
Fundamental Region of the Group of Fractional-C111/C222  
Linear Transformations

$$(1) \quad w = f(z) = \sum_{n=0}^{\infty} c_n z^n$$

regular within the circle  $|z| < 1$  and mapping this circle onto the regions of the family  $\{D\}$ . The given problem is reduced to the determination of the maximum of  $|f'(0)|$  in the class  $S_a(T)$ .

Theorem 1 : If the functional  $|f'(0)|^a$  has its maximum for  $w = f(z) \in S_a(T)$

then  $f$  maps the circle  $|z| < 1$  onto a region  $D$  with the following properties:

1)  $D$  contains  $w = c_0$  and is a simply connected fundamental region  $S_0$  of the group  $\{T\}$  with cuts. The boundary of  $D$  consists of finitely many analytic arcs being pairwise congruent with respect to  $\{T\}$ , and of piecewise analytic cuts beginning on the boundary of  $S_0$  and ending in the points  $a_1, \dots, a_m$  or in congruent points. 2) To every pair of congruent arcs of the boundary of  $S_0$  and to the simple arcs of the cuts there

Card 2/4

85944

On the Maximum of the Conformal Radius of the Fundamental Region of the Group of Fractional-Linear Transformations S/039/60/052/001/006/009XX C111/C222

there correspond two arcs of the same length on  $|z| = 1$  in consequence of  $w = f(z)$ .

Theorem 2 : The region D with the properties 1) and 2) of theorem 1 is unique.

Several exceptional cases are considered : 1)  $\{T\}$  consists of the transformations

$$w' = \frac{aw+b}{cw+d}, \quad ad - bc = 1, \quad a, b, c, d - \text{real, integral; } 2) \text{ as}$$

1) and besides  $b = 2b', c = 2c', b', c' - \text{integral; } 3) \{T\}$  consists of

$$w' = \frac{a\bar{w} + c}{c\bar{w} + a}, \quad a\bar{a} - c\bar{c} = 1, \quad a \text{ and } c - \text{complex integers. In all three cases}$$

the inverse function  $z = \varphi(w)$  of the extremal function is given explicitly ; the given formulas can be obtained from the general results of M.A. Lavrent'yev (Ref. 2). Finally, as a more complicated special case, the author considers groups whose fundamental regions are mappings of Riemannian surfaces (in this case there exists no simple automorphic

Card 3/4

On the Maximum of the Conformal Radius of the <sup>85944</sup> S/039/60/052/001/006/009 XX  
Fundamental Region of the Group of Fractional- C111/C222  
Linear Transformations

function having only one simple pole in the fundamental region.)  
The author mentions G.M. Goluzin and I.Ye. Bazilevich.  
There are 7 references : 6 Soviet and 1 English.

SUBMITTED: January 14, 1959

Card 4/4

S/044/62/000/G07/013/100  
C111/C333

AUTHOR: Gel'fer, S. A.

TITLE: Some extremal properties of multi-valent conformal mappings

PERIODICAL: Referativnyy zhurnal, Matematika, no. 7, 1962, 27-28, abstract 7B135. ("Issled. po sovrem. probl. teorii funktsiy kompleksn. peremennogo." M., Fizmatgiz, 1961, 351-359)

TEXT: Let  $E$  be the unit circle  $|z| < 1$ ; let  $K_z(r_0, R_0) \equiv K$  be the circular ring  $r_0 < |z| < R_0$ ,  $r_0 < 1 < R_0$ ; let  $z_0$  be a fixed point of  $E$  or  $K$ ;  $p \geq 1$ ,  $g \geq 0$ -integers;  $\alpha$ -- real number. Let  $R_p^{(g)}$  denote a closed Riemann surface with  $p$  sheets and genus  $g$ . The author gives a qualitative solution of the following extremal problem: The maximum of the functional

$$I_p = \operatorname{Re} \left[ e^{i\alpha} f(z_0) \right] \quad (1)$$

is determined, if  $f(z)$  varies in the class  $R_p^g$  in  $E$  or the class

Card 1/4

S/044/62/000/007/013/100  
C111/C333

Some extremal properties of . . .

$F_p \left( \frac{r_0}{R_0}, \frac{1}{P_1} \right) = F_p$  in  $K$ . Here  $F_p^g$  is the class of regular functions  $w = f(z) = z + c_2 z^2 + \dots + c_n z^n + \dots$  ( $f(z) \neq 0$  for  $z \neq 0$  in  $E$ , which map  $E$  one-to-one on the domain  $D$ ; the domains  $D$  lie on  $R_p^{(g)}$  and  $z = 0$  is thereby mapped on a fixed point  $O$  of  $R_p^{(g)}$  situated above  $w = 0$ .

Analogously, under the class

$F_p \left( \frac{r_0}{R_0}, \frac{1}{P_1} \right) = F_p$  the author understands the class of regular functions  $w = f(z)$ ,  $f(z) \neq 0$  in  $K$ , which map  $K$  one-to-one on twofold connected domains  $D$  on  $R_p^{(0)}$ , whereby  $z = 1$  is mapped on a fixed point  $P_1$  of  $R_p^{(0)}$  situated above the point  $w = 1$ . Both classes are normal in  $E$  and  $K$ , respectively, this guarantees the existence of extremal mapping functions of the extremal problem being considered in the same class.

Card 2/7



S/044/62/000/007/013/100  
C111/C333

Some extremal properties of . . .

To solve the given problem the author extends the variation method by G. M. Goluzin (Goluzin, G. M., Geometricheskaya teoriya funktsiy kompleksnogo peremennogo [Geometric function theory] M.-L., 1952) to the mentioned function classes by constructing the corresponding variation formulas for these classes. The application of these variation formulas to the extremal problem being examined leads to differential equations in E and K, respectively, for the sought extremal mapping functions, thereby permitting the character of the mappings effected by these functions to be clarified. The following theorems result:

Theorem 1.1: The function  $f(z) \in F_p^g$ , for which (1) attains a maximum, maps the circle  $|z| < 1$  on the whole surface  $R_p^{(g)}$  with one cut consisting of a finite number of analytical arcs. These arcs contain all points of  $R_p^{(g)}$  which are situated over the points  $w = 0$  and  $w = \infty$  of the w-plane with the exception of the point 0.

Theorem 2.1: The function  $f(z) \in F_p$  for which (1) attains a maximum maps K on a domain D which originates from  $R_p^{(0)}$  with the help of two  
Card 3/4

Some extremal properties of . . .

S/044/62/000/007/013/100  
C111/C333

cuts consisting of a finite number of analytical arcs. These arcs contain all points of  $R_p^{(0)}$  which are situated over the points  $w = 0$  and  $w = \infty$  of the  $w$ -plane. ✓

Further, a differential equation is obtained for the case  $z \in E$ ,  $f \in F_p^E$ , the integral curves of which are analytical arcs which form the boundary cuts of the extremal domain from theorem 1.1.

[Abstracter's note: Complete translation.]

Card 4/4

34736

3/020/62/142/003/001/027  
C 111/C333

16.3000

AUTHOR: Gel'fer, S.A.

TITLE: Extension of the Goluzin-Schiffer variational method to multiply connected regions

PERIODICAL: Akademiya nauk SSSR. Doklady, v. 142, no. 3, 1962, 503-506

TEXT: Let the  $n$ -fold connected domain  $D$  be the ring  $1 < |z| < R$  with  $n - 2$  circular holes. The class  $P$  is assumed to consist of the functions  $w = f(z)$ ,  $f(z) \neq 0$  regular and schlicht in  $D$  for  $z \in D$ ,  $f(z_0) = 1$  for a fixed  $z_0 \in (1, R)$  with the property that, if the curve  $l_1 \subset D$  is contained in the curve  $l_2 \subset D$ ,  $f|_{l_1}$  is contained in  $f|_{l_2}$ . Let  $\phi[f]$  be a real functional defined for all analytic functions in  $D$  which satisfies the relation

$$\phi[f + \varepsilon g] = \phi[f] + \operatorname{Re}\{\varepsilon \psi[f, g]\} - O(\varepsilon^2) \quad (1)$$

from M. Schiffer (Ref. 1: Proc. Internat. Congr. Math., 1958, Cambridge, 1960), where  $\psi[f, g]$  is a linear complex functional in  $g(z)$ .

Card 1/4

Extension of the Goluzin-Schiffer ...

S/020/62/142/003/001/027  
C111/C333

The following problem is considered : Determine the functions of class  $P$  for which  $\Phi[f]$  becomes maximum.  
In order to solve the problem the author goes back to a result of V.D. Yerokhin (Ref. 2 : DAN, 120, no. 4, 689 (1953); 127, no. 6, 1155 (1959)) and represents the sought extremum function in the form

$$f(z) = f_n f_{n-1} \dots f_1(z), \quad z \in D, \quad (2)$$

where  $f_1, f_2, \dots, f_n$  are unique and conformal in the domains  $D(L_1), f_1 D(f_1 L_2), \dots, f_{n-1} \dots f_1 D(f_{n-1} \dots f_1 L_n)$ ; here  $D(L)$  is a simply connected domain containing  $D$  and having the boundary  $L$ ;  $L_1: |z| < 1$ ,  $L_n: |z| < R$ ;  $L_2, \dots, L_{n-1}$  the other  $n - 2$  circles of the boundary of  $D$ . The component  $t_1 = f_1(z)$  is chosen such that the conditions  $f_1(\infty) = \infty$ ,  $f_1(z) \neq 0$ ,  $f_1(z_0) = t_1^0$  are satisfied, where  $t_1^0$  is determined from  $f_n f_{n-1} \dots f_2(t_1^0) = 1$ . Then the variation formula of G.M. Goluzin

Card 2/4